


## Patna Sahib (GEN)

Bihar

No. of Electors (2019): 21,46,008

### Intersecting Districts

Patna



## What this Factsheet Contains

This Factsheet is based on our interactive **NFHS Policy Tracker for Parliamentary Constituencies (PCs)**—part of our **India Policy Insights** initiative. It provides performance measures for 93 indicators and a side-by-side comparison with national and state averages for each PC. It also shows the net change between 2016 and 2021 for 86 indicators to highlight performance gains and losses. The Factsheet additionally lists the intersecting districts for each PC and links each indicator to a current GOI program, thereby readily providing a policy and program context for its meaningful use. See: <https://geographicinsights.iq.harvard.edu/nfhs-tracker-pc>

### Indicator Summary (2021)

Number of Indicators better than All-India average



Number of Indicators better than State average




### Indicator Change Summary (Between 2016 and 2021)

Ranking based on Percentage of Indicators showing Any Improvement (out of 543 PCs)



● Highest Improvement | ● Improvement | ● Worsened | ● Extremely Worsened



#### Top 15 Indicators showing Improvement

- Alcohol consumption (Men)
- Menstrual hygiene
- Iron folic acid for 180 days
- Receiving solid/semi-solid food (6-8 months)
- Tobacco use (Men)
- Children with diarrhoea taken to health facility
- Any contraceptive method
- Mother and Child Protection card received
- Stunting
- Exclusive breastfeeding (within 6 months)
- Children with fever or ARI taken to health facility
- Birth registration
- Underweight
- Population in household with electricity
- Unmet need for spacing

#### Top 15 Indicators showing Worsening

- Anemia (Pregnant women)
- Anemia (Children)
- High or very high or taking medicine (Men)
- Full vaccination
- Anemia (Non-pregnant women)
- High or very high or taking medicine (Women)
- Anemia (All women)
- Full vaccination (Source from card only)
- 3 doses of polio vaccine
- Children with diarrhoea who received zinc
- Current users ever told about side effects
- Third or higher order birth
- Caesarean section in private sector
- Last birth protected against neonatal tetanus
- At least 4 antenatal care visits

For more details, see the complete listing of Improving or Worsening Indicators (pp. 2-4).

Indicator %	Government of India Programs	India 2021	State 2021	PC 2021	PC Change 2021-2016	
<b>Population and Household Profile</b>						
Birth registration	RBDA	89.1	75.6	<b>71.4</b>	3.8	●
Female school attendance	RTE, NEP, SSA, KGBV, DGAG, NSIG, SVSGC, BBBP, SWAYAM	71.8	61.1	<b>73</b>	0.6	●
Health insurance coverage	PMJAY, PMSBY, AABY	41	14.6	<b>10.1</b>	2.4	●
Improved sanitation	SBM	70.2	49.4	<b>66.8</b>	0.9	●
Population in household with electricity	Saubhagya, DDUGJY, KUSUM, Ujjwala-DISCOM, IPDS, SCM, JNNURM, WISTS, NSM, PMAY-U, JNNURM, AMRUT, MPLADS	96.8	96.3	<b>99.4</b>	3.2	●
Clean fuel for cooking	PMUY	58.6	37.8	<b>71.3</b>	-0.6	●
Improved water	JJM, ABY, Amrit sarovar, Jaldoot, PMAY-G, SBM-G	95.9	99.2	<b>98.7</b>	-0.5	●
Population below 15 years	RKSK, RBSK	26.5	36.4	<b>31.6</b>	-0.5	●
Iodized salt	NIDDCP, Fortification	94.3	93.3	<b>94.4</b>	-4.9	●
Death registration	RBDA, MDSR	70.8	37.1	<b>40.9</b>		
<b>Characteristics of Women</b>						
10 or more years of schooling (Women)	RUSA, ATL, RAA, DIKSHA, NEP	41	28.8	<b>49.5</b>	1.2	●
Literate women	NILP	71.5	57.8	<b>70.9</b>	-0.6	●
<b>Marriage and Fertility</b>						
Menstrual hygiene	SPMH, AFHC, AHS, PEP, RKSK, SBM-G	77.6	58.8	<b>80.2</b>	16.5	●
Child marriage (Women)	PCM	23.3	40.8	<b>26.6</b>	0.1	●
Adolescent pregnancy	POCSO, MTP	6.8	11	<b>9.5</b>	1.5	●
Third or higher order birth	ESB, HDC, MPV, MTP, NPP	26.5		<b>42.7</b>	7.3	●
<b>Current Use of Family Planning Methods</b>						
Pill	NPP, MPV, Chayya, Mala-N, NRHM	5.4	2	<b>2</b>	1.4	●
Any contraceptive method	NPP, MPV, NRHM, FPIS, ECS, RPF-PPIUCD, Antara, ESB, HDC	66.7	55.8	<b>49.6</b>	8.3	●
Any modern method	NPP, MPV, NRHM, FPIS, ECS, RPF-PPIUCD, Antara, ESB, HDC	56.4	44.4	<b>42.2</b>	2.6	●
Condom	NPP, MPV, HDC, NRHM	9.5	4	<b>4.1</b>	2.3	●
Male sterilization	MPV, NSV, FPIS, NRHM	0.3	0.1	<b>0.1</b>	0	●
Female sterilization	MPV, FPIS, NRHM	37.9	34.8	<b>31.3</b>	-2.3	●
IUD/PPIUD	NPP, MPV, RPF-PPIUCD, NRHM	2.1	0.8	<b>0.9</b>	-0.2	●
Injectables	NPP, MPV, Antara, NRHM	0.6	1.1	<b>0.5</b>		
<b>Unmet Need for Family Planning</b>						
Unmet need for spacing	NPP, MPV, ESB, NRHM, FPIS, ECS, RPF-PPIUCD, Antara, HDC	4	6.1	<b>4.7</b>	-2.6	●
Total unmet need	NPP, MPV, RPF-PPIUCD, Antara, ESB, HDC, NRHM, FPIS, ECS	9.4	13.6	<b>15.4</b>	-1.4	●
<b>Quality of Family Planning Services</b>						
Interaction of health worker	MPV, NRHM, ESB, JSSK	23.9	20.2	<b>24.7</b>	-1.6	●
Current users ever told about side effects	NPP, MPV, NRHM	62.4	49.9	<b>45.5</b>	-7.5	●
<b>Maternal and Child Health</b>						
Iron folic acid for 100 days	AMB, POSHAN, NAP, WIFSP, RBSK, TBSY	44.1	18	<b>20.5</b>	1.7	●
Iron folic acid for 180 days	AMB, POSHAN, NAP, WIFSP, RBSK, TBSY	26	9.2	<b>13.9</b>	11.9	●

Indicator %	Government of India Programs	India 2021	State 2021	PC 2021	PC Change 2021-2016	
Mother and Child Protection card received	None	95.9	89.5	<b>92.2</b>	8.2	●
Antenatal check-up in the first trimester	PMSMA, PMMVY, JSSK	70	52.9	<b>54.9</b>	-3.5	●
Mothers received postnatal care	PMSMA, PMMVY, MCC, JSSK	78	57.3	<b>59.8</b>	-4.4	●
Postnatal check by skilled health workers	JSSK, SMA, MCC, ICDS, NSSK	11.5	59.3	<b>4.4</b>	-0.7	●
At least 4 antenatal care visits	PMSMA, PMMVY, JSSK	58.5	25.2	<b>18.3</b>	-6.7	●
Last birth protected against neonatal tetanus	PMMVY, PMSMA, SMA, ICDS, JSSK, LQI	92	89.5	<b>87.9</b>	-6.8	●
<b>Delivery Care</b>						
Home delivery by skilled health personnel	BMBP, HBNC, SMB, PMMVY, ICDS-ANM	3.2	6.1	<b>1.9</b>	0.5	●
In public facility	JSY, JSSK, LQI, PMMVY, PMSMA, MCC	61.9	56.9	<b>46</b>	1.6	●
Institutional births	JSY, JSSK, LQI, PMMVY, PMSMA, MCC	88.6	76.2	<b>93</b>	1.6	●
Caesarean section delivery	JSY, JSSK, LQI	21.5	9.7	<b>21</b>	3.6	●
Caesarean section in private sector	JSY, JSSK, LQI	47.5	39.6	<b>45.5</b>	7.3	●
Caesarean section in public sector	JSY, JSSK, LQI	14.3	3.6	<b>5.6</b>	2	●
Births by skilled health personnel	JSY, JSSK, LQI, HBNC	89.4	79	<b>84.8</b>	-6.1	●
<b>Child Vaccinations and Vit A Supplements</b>						
Vitamin A dose in the last 6 months	NVAPP	71.2	56	<b>60.4</b>	1.1	●
3 doses of penta or DPT vaccine	IMI	86.7	85	<b>86.5</b>	-2.1	●
3 doses of penta or hepatitis B vaccine	IMI	83.9	82.3	<b>75</b>	-3.3	●
First dose of measles-containing vaccine	IMI	87.9	85.7	<b>84.3</b>	-2.5	●
3 doses of polio vaccine	IMI, PPP	80.5	75.5	<b>67.4</b>	-10.4	●
Full vaccination	IMI, PPP	76.6	71	<b>58.3</b>	-13.4	●
Full vaccination (Source from card only)	IMI, PPP	74.3	82.7	<b>75.9</b>	-10.5	●
3 doses of rotavirus vaccine	IMI	36.4	3.4	<b>6.4</b>		
Second dose of measles-containing vaccine	IMI	58.6	29.9	<b>40.1</b>		
<b>Treatment of Childhood Diseases</b>						
Children with diarrhoea taken to health facility	RBSK, IDCF	76.2	64.7	<b>72.6</b>	9.5	●
Children with fever or ARI taken to health facility	NTEP, RBSK, SANPS	69	69.3	<b>70.5</b>	4.6	●
Prevalence of ARI	NTEP, RBSK, SANPS, RM-ARI	2.8	3.5	<b>0.7</b>	0	●
Children with diarrhoea who received ORS	RBSK, IDCF	60.6	58.2	<b>52.7</b>	-1	●
Children with diarrhoea who received zinc	RBSK, IDCF	30.5	25.6	<b>23.5</b>	-9.4	●
Prevalence of diarrhoea	RBSK, IDCF	7.3	13.7	<b>8</b>	4.1	●
<b>Child Feeding and Nutritional Status</b>						
Stunting	POSHAN, ICDS, AK, NRC, RBSK, BPKK	35.5	42.9	<b>34</b>	-7.1	●
Exclusive breastfeeding (within 6 months)	MAA, LMC	63.7	58.9	<b>44.8</b>	6.4	●
Breastfed within one hour of birth	MAA, LMC, JSSK	41.8	31.1	<b>33.5</b>	-5.6	●
Overweight	POSHAN	3.4	2.4	<b>1.3</b>	0.3	●

Indicator %	Government of India Programs	India 2021	State 2021	PC 2021	PC Change 2021-2016	
Receiving solid/semi-solid food (6-8 months)	POSHAN, ICDS	45.9	39	<b>41.2</b>	11.4	●
Adequate diet (All children - 6-23 months)	NFSA, TPDS, POSHAN, WBNP, AK, RKVY, PMMVY, AAY	11	10.9	<b>6.2</b>	2.1	●
Adequate diet (Breastfeeding - 6-23 months)	MAA, LMC, NFSA, POSHAN	10.8	10.8	<b>6</b>	2.4	●
Severe wasting	POSHAN, ICDS, AK, NRC, RBSK, BPKK	7.7	8.8	<b>9</b>	-1.1	●
Underweight	POSHAN, ICDS, AK, NRC, RBSK, BPKK	32.1	41	<b>38.4</b>	-3.6	●
Wasting	POSHAN, ICDS, AK, NRC, RBSK, BPKK	19.3	22.9	<b>25.8</b>	-1.1	●
Adequate diet (Non-breastfeeding - 6-23 months)	NFSA, TPDS, POSHAN, WBNP, AK, RKVY, PMMVY, AAY	12.3	11.5	<b>5.8</b>	-1.7	●
<b>Nutritional Status of Women</b>						
Overweight or obese (Women)	IHIP, PMSSY, NPCDCS	24	15.9	<b>21.5</b>	-0.6	●
BMI below normal (Women)	AK, NRC, BPKK, NFSA, TPDS, POSHAN	18.7	25.6	<b>21.4</b>	0	●
Risky waist-to-hip ratio (Women)	IHIP, PMSSY, NPCDCS	56.7	60.3	<b>68.7</b>		
<b>Anemia among Children and Women</b>						
Anemia (All adolescent women)	AMB, NAP, WIFSP, RSKS, TBSY	59.1	65.7	<b>62.1</b>	6.4	●
Anemia (All women)	AMB, NAP, WIFSP	57	63.5	<b>66.6</b>	11.9	●
Anemia (Children)	AMB, POSHAN, NAP, WIFSP, RBSK, TBSY	67.1	69.4	<b>65.7</b>	14.9	●
Anemia (Non-pregnant women)	AMB, NAP, WIFSP	57.2	63.6	<b>69</b>	13.3	●
Anemia (Pregnant women)	AMB, NAP, WIFSP, PMSMA, JSSK	52.2	63.1	<b>61.9</b>	17.8	●
<b>Blood Sugar Level among Adults</b>						
High (141-160 mg/dl) (Men)	NPCDCS, AMRIT, eAushidhi, IHIP, PMSSY, ABWC	7.3	8.3	<b>9.1</b>	6.2	●
High (141-160 mg/dl) (Women)	NPCDCS, AMRIT, eAushidhi, IHIP, PMSSY, ABWC	6.1	6.4	<b>7.1</b>	5.1	●
High or very high or taking medicine (Men)	NPCDCS, AMRIT, eAushidhi, IHIP, PMSSY, ABWC	15.6	16.2	<b>20.8</b>	13.5	●
High or very high or taking medicine (Women)	NPCDCS, AMRIT, eAushidhi, IHIP, PMSSY, ABWC	13.5	12.7	<b>16.8</b>	12.3	●
Very high (>160 mg/dl) (Men)	NPCDCS, AMRIT, eAushidhi, IHIP, PMSSY, ABWC	7.1	7	<b>9.2</b>	5.8	●
Very high (>160 mg/dl) (Women)	NPCDCS, AMRIT, eAushidhi, IHIP, PMSSY, ABWC	6.3	5.4	<b>5.8</b>	3.9	●
<b>Hypertension among Adults</b>						
Mild blood pressure (Men)	NPCDCS, HIHC, AMRIT, eAushidhi, IHIP, PMSSY, ABWC	15.7	11.2	<b>10.4</b>	2.5	●
Mild blood pressure (Women)	NPCDCS, HIHC, AMRIT, eAushidhi, IHIP, PMSSY, ABWC	12.4	8.7	<b>7.5</b>	3.7	●
Moderate or severe blood pressure (Men)	NPCDCS, HIHC, AMRIT, eAushidhi, IHIP, PMSSY, ABWC	5.7	4.3	<b>2.5</b>	0.4	●
Moderate or severe blood pressure (Women)	NPCDCS, HIHC, AMRIT, eAushidhi, IHIP, PMSSY, ABWC	5.2	3.6	<b>2.5</b>	1.2	●
Blood Pressure - High/Medication (Men)	NPCDCS, HIHC, AMRIT, eAushidhi, IHIP, PMSSY, ABWC	24	18.4	<b>15.6</b>		
Blood Pressure - High/Medication (Women)	NPCDCS, HIHC, AMRIT, eAushidhi, IHIP, PMSSY, ABWC	21.3	15.9	<b>16</b>		
<b>Tobacco and Alcohol Use among Adults</b>						
Alcohol consumption (Men)	IRCA, NPCDCS	18.8	15.5	<b>21.5</b>	-19.9	●
Tobacco use (Men)	NTCP, COTPA, PEC	38	48.8	<b>38.9</b>	-9.8	●
Alcohol consumption (Women)	IRCA, NPCDCS	1.3	0.4	<b>0</b>	0	●
Tobacco use (Women)	NTCP, COTPA, PEC	8.9	5	<b>0.2</b>	-0.5	●

